



Volume 1 | Issue 1 | Jan - Jun 2023

ISSN: 2984-9896

e-ISSN: 3116-5230

Website: <https://csu.org.ph/jpas>

Email: csujpas@gmail.com



Burn Wound Healing Potential of Paragis (*Eleusine indica*) Leaf Ethanolic Extract Ointment on Sprague Dawley (*Rattus norvegicus*) Rats

Celso L. Baccay Jr., Grescen Alelee B. Glinoga, Donna Mae M. Lumaban and Madel B. Tanguilan
Cagayan State University – Carig Campus, Carig Sur, Tuguegarao City, Cagayan, Philippines

Corresponding Author: Madel B. Tanguilan ✉ mtancsu@gmail.com

DOI: [10.5281/zenodo.20710590](https://doi.org/10.5281/zenodo.20710590)

Published: June 30, 2023

ABSTRACT

This study aimed to investigate the burn wound healing potential of the ethanolic extract and 10% ointment from Paragis (*Eleusine indica*) leaves. The research utilized the Complete Randomized Design in evaluating the percent wound contraction and epithelialization period (days) of burn wound inflicted on the test animals, Sprague Dawley rats. The 10% ointment prepared from the ethanolic extract of Paragis (*E. indica*) leaves was found to be non-cytotoxic in the acute dermal toxicity test at the different dosage levels: 2000 mg/kg (limit test), 1000 mg/kg, 500 mg/kg, and 200 mg/kg. The ointment (Treatment 1) produced with ethanolic leaf extract of Paragis, showed the highest percentage (contraction) for it reached 100% within 16 days which was the last day of monitoring, followed by Treatment 2 (Paragis leaf ethanolic extract), because it reached 93.32% wound contraction at the expected day of contraction (16th day). The positive control ranked third with 84.98% wound contraction at day 16, followed by negative control with 66.62% wound contraction.

Keywords: Burn wound healing, Ethanolic extract, Sprague Dawley rats



INTRODUCTION

Wound healing is a crucial mechanism for the maintenance of homeostasis. In this regard, the skin must sustain a “robust and effective repair mechanism to ensure restoration of tissues and regeneration of injured skin or tissues. Cutaneous wound healing emanates as the “skin repairs itself” after exposure to injury due to “surgery, trauma, and burns”. Wound healing occurs as a systematic flow of processes from “inflammation, angiogenesis, proliferation and synthesis of collagen for final healing” (Hemamalini et al. as cited in Ahmed et al., 2016). Paragis belongs to the Class Magnoliopsida, Family Poaceae, with the binomial name *Eleusine indica*. It is a weed that grows abundantly in warm areas, particularly “along riverbanks, roads, and settled areas throughout the Philippines”. In the Philippines, the decoction of leaves serves as a diuretic (Surigao del Sur) and a treatment for kidney diseases and arthritis (Sablan, Benguet Province) whereas dried leaves and stems are burned as an insect repellent (Porac, Pampanga). The plant is considered to possess folkloric medicinal value in different countries. It is recognized as a “diuretic, antihelmintic, diaphoretic, and febrifuge”. Paragis is also used against “diarrhea, dysentery, epilepsy, and intestinal occlusion” (Cameroon), “treatment of infertility in females,” muscle sprains (Sri Lanka), hypertension (Myanmar), “treatment of prolapsed uterus” (Bangladesh), liver disorders, and snake-bite (India) (Bangladesh Ethnobotany Online Database; Biswas et al.; Mabeku; Obico & Ragragio; Upasani et al; Wai et al., as cited in Stuart, 2013). These traditional uses of Paragis should be probed further.

Several studies provide evidence of the medicinal efficacy of Paragis. This plant is noted to possess “antiviral, antiplasmodial, antidiabetic, and antibacterial properties. It was also investigated due to its “anticancer potential” (Aye et al, Morah & Otuk as cited in Nas et al, 2020).

Evidently, Paragis is one plant that deserves further scientific inquiry. Thus, this study will add to the existing

body of knowledge on the wound healing capacity of an ointment from the ethanolic leaf extract of Paragis.

Objectives of the Study

This study aims to assess the wound healing potential of Paragis (*Eleusine indica*) leaf ethanolic extract ointment. It intends to examine the physical characteristics of the extract, perform physico-pharmaceutical evaluations of the ointment, determine acute dermal toxicity, and assess wound healing by measuring wound contraction and epithelialization. Additionally, the research seeks to establish whether Paragis ethanolic extract and Paragis ointment exhibit significant differences in wound healing compared to a positive control group.

MATERIALS AND METHODS

Research Design

The experiment was performed using a Complete Randomized Design (CRD) with a total of twenty-four (24) Sprague Dawley rats, which were distributed in the following order: four (4) rats for the acute dermal toxicity test (200 mg/kg, 500 mg/kg, 1000 mg/kg, and 2000 mg/kg ointments) and twenty (20) with five (5) replicates each for the four (4) treatments (negative control, positive control (0.2% Nitrofurazone), 10% ointment, and ethanolic extract) for the burn wound model.

- T1 – 10% ointment (from Paragis leaves ethanolic extract)
- T2 – Ethanolic extract of the Paragis leaves
- T3 – positive control (0.2% Nitrofurazone)
- T4 – negative control (untreated).

Sampling Technique

A group of healthy Sprague Dawley rats, each weighing between 200 to 250 grams, were randomly chosen as the study subjects.

Locale of the study

The Paragis leaves used in this study were gathered from the outskirts of Gosi Sur, situated in Tuguegarao City, Cagayan. The experimental procedures were

meticulously carried out within the confines of a controlled laboratory environment.

Research Instruments

The experimental setup for this study adhered to ethical standards and guidelines, with an animal permit secured from the Institutional Animal Care and Use Committee (IACUC). To ensure accurate and precise results, various chemicals and specialized equipment were utilized for plant extraction and ointment preparation. Additionally, Lidocaine was employed for anesthesia purposes, while normal saline played a crucial role in wound cleaning. To measure wound areas with precision, a transparent ruler was employed as a valuable tool throughout the experiment.

Data Gathering Procedure

Plant Collection and Authentication

Fresh, mature leaves of Paragis, approximately five kilograms (5 kg) were randomly collected by the researchers from the border of Gosi Sur, Tuguegarao City, Cagayan. It was authenticated by the Department of Agriculture Region II. The duration of collection lasted for about three (3) days. The leaves were washed initially with tap water and then distilled water to ensure the removal of soil and other impurities that may affect the result of the experiment. The washed leaves were cut into smaller strips and allowed to air dry until a constant weight was observed.

Preparation of the Plant Extract and Ethanolic Extract

The method of ethanol extraction was adapted from the procedure presented by Guevara (2005). The air-dried leaves of Paragis were subjected to ethanolic extraction. About 300g of sample was soaked in 3000 ml (1:10 g/ml ratio) of 70% ethanol for 48 hours. The filtrate was collected after filtration of the residue using a Whatman No. 1 filter paper and then concentrated by Rotary evaporator. A flame test was conducted to ascertain the absence of ethanol in the extract of Paragis leaves. The extracts were then packed in a clean, dry, and tightly

sealed bottle and refrigerated until further use in the bum wound model.

Physical Characteristics of Paragis Leaves Ethanolic Extract

Physical characteristics of Paragis extract such as its color and odor were evaluated by visual and olfactory observation. The color and odor of the Paragis leaf ethanolic extract was observed and noted. Production of 10% Ointment from the Paragis Leaves Ethanolic Extract (Fusion Method Adapted from Rao Kodati et al., 2011). In the preparation of 10% ointment, two grams (2g) Wool fat (Lanolin), two grams (2g) of Hard Paraffin, two grams (2g) of Cetostearyl alcohol, and thirty-four grams (34 g) White Soft Paraffin were used. Each ingredient was mixed, the mixture was heated gently with constant stirring to achieve the desired consistency, and then the preparation was cooled. Four milliliters (4 mL) of Ethanolic Extract of Paragis leaves was slowly added to the melted ingredients and then stirred thoroughly to obtain a homogeneous mixture. The ointment was then packed in a wide mouth container.

Physico-pharmaceutical Evaluation of the 10% Ointment

The pH was determined using a pH meter. External characteristics such as color, odor, smoothness, grittiness, and homogeneity were determined by visual inspection and tested for appearance with no lumps.

Experimental Protocol on the Acute Dermal Toxicity and Wound Healing Activity

Acclimatization of Animals

The researchers secured an Animal Permit from Institutional Animal Care and Use Committee (IACUC). Healthy Sprague Dawley rats were randomly selected for this experiment. As noted by Demilew et al. (2018), experimental animals weighed 200 -250g. The animals were kept in individual cages where they had free access to diet and water for two (2) weeks for acclimatization to the laboratory conditions.

Grouping and Dosing of Animals

Grouping and dosing of the animal specimen were done as follows: four (4) treatment groups, each with five (5) rats were used for the burn wound model. Animals in Treatment I received the 10% ointment from the ethanolic extract of Paragis leaves. Those in Treatment II were treated with Paragis leaf ethanolic extract. Animals in Treatment III received 0.2% Nitrofurazone (Positive Control) while those in treatment 4 (Negative Control) were left untreated after the burn infliction. The remaining four (4) Sprague Dawley rats were utilized for the acute dermal toxicity test. All the conducted experiments conformed to the international accepted guideline for laboratory animal use and care published by the National Academies Press (as cited in Demilew et al., 2018).

Acute Dermal Toxicity Test (Adapted from OECD November 22, 2017)

One rat was assigned per dose. The test substance was applied to the skin (not less than 10 percent of the body surface area) in graduated doses on each experimental animal. At least three dose levels were used, excluding the limit test, appropriately spaced to produce a dose-response curve. A limit test of at least 2000 mg/kg was made. The observation period was at least fourteen (14) days. During the first day, the animals were observed for five (5) hours and then the observations were made daily.

Burn Wound Model (Adapted from Subalakshmi, 2014)

Hairs from the predetermined area (refer to figure 1) for burn wound infliction at the back of the Sprague Dawley rats were removed with a previously sanitized razor. The skin lateral to the spinal groove was shaved, cleaned with cool distilled water, and then dried. The burn infliction done by Subalakshmi et al. (2014) was adopted in this study. The grouped rats were injected with Lidocaine (anesthesia) in the dorsal region of their body. A cylindrical metal rod of about twelve (12) mm in diameter, which was heated for 30 seconds with a blow torch, was pressed to the shaved portion of the rat for six

(6) seconds. All wounds were cleaned with normal saline suggested by Khoo et al. (2010). The animals were then placed individually in their own cages after the infliction of the burn wounds and for recovery from the anesthesia.



Figure 1. Site of burn wound infliction in

The animals were monitored immediately postoperatively for spontaneous breathing efforts and movement. After surgery, each animal was housed in an individual cage in a room and fed with standard rat diet and water. All wounds were cleaned with normal saline, and treatment was reapplied every three days.

Measurement of Wound Contraction (Adapted from Nagar et.al., 2016)

The burn wound healing potential was monitored by measuring the wound mean area and computing the percentage (%) of wound contraction among the four (4) treatments. The monitoring and assessment protocol suggested by Nagar et al. (2016) was used, except that the wound contraction evaluation was done on days 1, 4, 7, 10, 13, and 16 using a transparent ruler instead of a tracing paper. In case of infection occurrence, the monitoring was extended for two (2) more days (Subalakshmi et al. as cited in Demilew et al, 2018). Measurements of wound area were done on 1 mm² scale. The surface area recorded on each rat was used to calculate the percent wound contraction, using the number of days (eighteen days) and taking the initial size of the wound as 100%, with the following formula:

$$\begin{aligned} & \% \text{ wound contraction} \\ & = (\text{wound area on 1}^{\text{st}} \text{ day} - \text{wound area on} \\ & \text{day (n)}) / \text{Wound area on 1}^{\text{st}} \text{ day} \times 100 \end{aligned}$$

Epithelialization Period

The falling of the scab, without traces of wound, served as the basis for the epithelialization period. This event was noted as the “endpoint of complete epithelialization and the days required for this were taken as period of epithelialization” (Subalakshmi et al. as cited in Demilew et al., 2018).

Analysis of the Data/ Statistical treatment

The independent variable in the study was the type of treatment negative control (untreated), positive control (0.2% Nitrofurazone), ethanolic extract, and 10% ointment from the ethanolic extract of the Paragis leaves. The dependent variables, analyzed separately, were the percent of wound contraction and epithelialization period (days). The data on percent wound contraction on days 10, 13, and 16 were statistically analyzed. The data (percent wound contraction and epithelialization period) were analyzed using One-way Analysis of Variance (ANOVA). At a 95% confidence level (significance level of $\alpha = 0.05$), a p-value < 0.05 was used to reject the null hypothesis, which indicated a statistically significant difference in the efficacy of the test samples (10% ointment and ethanolic extract) against the positive control (0.2% Nitrofurazone). In case of rejection of the null hypothesis, further tests were done (Single Factor ANOVA Pairwise Comparison) to identify the treatments/groups that elicited the significant difference.

RESULTS AND DISCUSSION

Table 1. Physical Evaluation of Paragis leaf extract.

Physical Evaluation		
	Standard	Observation
Color	Dark green- brown	Yellowish- brown
Odor	Aromatic	Dried leaves tea

Table 1 shows the physical characteristics of Paragis leaf extract as to its color and odor as observed by the researchers and other respondents. The result showed that the color yellowish-brown and dried tea leaves odor was similar to the standard data of observation by V.P.S. Bhadauria, Varsha, Gupta et al., (2019).

Table 2. Physio-pharmaceutical evaluation of the ointment.

	Standard	Observation
Ph	5-7	Acidic (6)
Odor	Aromatic	Smells like wool fat
Smoothness	Smooth	Smooth
Grittiness	Free from grittiness	Non-gritty
Homogeneity and appearance with no lumps	Smooth Free from grittiness Homogeneous	Homogeneous with no lumps

Table 2 shows the physico-pharmaceutical evaluation of the ointment fused with Paragis extract. The pH of the ointment was tested acidic with a pH of 6 and conformed to the standard pH as stated by Baravkar (2011). The odor was similar to wool fat odor due to its wool fat component. The ointment was homogenous without lumps, thus making it non-gritty, which is comparable to other prepared ointments in the study conducted by Baravkar (2011). According to Dr. Vijendra Nalamothu (2015), the importance of conducting the physico-pharmaceutical evaluation of ointment is that in the formulation components, a simple change in properties, such as pH, viscosity, the relative amounts of leaf extract, water, surfactants, stabilizers, droplet size, ionic nature, or the method of preparation, can often influence skin absorption and efficacy.



Figure 2. Acute Dermal Toxicity of Ointment adopted from OECD 2017. (a) 2000 mg/kg; (b) 1000 mg/kg; (c) 500 mg/kg and (d) 200 mg/kg.

The study also conducted dermal toxicity of the ointment with different levels of doses. The test served as a basis for establishing the dosage of Treatments, and it also provided information on dermal adsorption and the mode of toxic action of substance.

Figure 2 shows the result on dermal toxicity. It clearly shows that the ointment was not toxic for the test animals and did not show severe and enduring signs of distress and pain as it was the principle of the tests. A limit test was made which was 2000 mg/kg with following the three levels of doses (1000, 500, and 200), respectively, based on OECD (1987 updated November 22, 2017). As regards for the formula for getting the amount of dosage, the researchers converted the mg/kg to g/g depending on the weight of the rat.




Figure 2. Acute Dermal Toxicity of Ointment adopted from OECD 2017. (a) 2000 mg/kg; (b) 1000 ng/kg; (c) 500 mg/kg and (d) 200 mg/kg.

In evaluating the wound healing process through percent wound contraction, the researchers used the protocol from Nagar, HK. et al, (2016). Figure 8 shows the evaluation of the wound healing process through percent wound contraction. As observed, the ointment (treatment 1) produced with ethanolic leaf extract of Paragis, showed the highest percentage (contraction) for it reached 100% within 16 days, which was the last day of monitoring, followed by Treatment 2 (Paragis leaf ethanolic extract), because it reached 93.32% wound contraction at the expected day of contraction (16th day). The positive control ranked third with 84.98% wound contraction on day 16, followed by negative control with 66.62% wound contraction.

The trend for percent wound contraction was T1 > T2 > T3 > T4 (in which T1 had the fastest rate of wound repair as seen in figure 8). For the falling off of scabs, the trend was: T1 > T2 > T3 > T4 (in which T1 or 10% ointment had the

lowest or fastest period of epithelialization). The wound healing was evident in Appendix B in which replicates 2, 3, and 5 for T1 (10% ointment) and replicates 1, 4, and 5 for T2 (ethanolic extract) were completely cured on Day 13. At Day 16, all replicates for T1 (10% ointment); 1, 3, 4, and 5 for T2 (ethanolic extract); 1,2, and 5 for T3 (0.2% Nitrofurazone), and 3 for T4 (untreated) were totally cured. The data was analyzed using one-way ANOVA. The efficacies were obtained by comparing the time of wound contraction using the four Treatments. The values were converted into percentages using the mean index, to show significant differences among all treatments. Treatment means that are significant, were compared using Least Significant Differences (LSD). Based on the analyzed data shown in table 3, Treatment 1 and Treatment 2 are not significantly different from Treatment 3. Treatments 3 and 4 also showed that they are significantly different at 0.05 level of significance.

Based on the results shown in table 3, treatments with the highest mean value have the greater bum wound healing potential. The treatment 1 (Ointment) ranked the highest with 100% wound contraction followed by Treatment II (Ethanolic extract), having 93.32%, Treatment II (Positive Control) with 84.98%, and Treatment IV (Untreated) with 66.62%, respectively.

LSD results suggest that Treatments I, II and TIII are not significantly different in terms of wound healing activity. Treatment III and Treatment IV are not significantly different.

Table 3. Mean of each treatment.

Treatment	Mean (%)	Rank
1 Ointment	100.00 ^a	1
2 Extract	93.32 ^a	2
3 Positive	84.98 ^{ab}	3
4 Negative	66.62 ^b	4
Total	86.23	

*Any two means having a common letter(s) are not significantly different, otherwise they are significantly different at 5% level of significance.

In evaluating the Period of Epithelialization, the falling of scabs was determined by researchers as cited in Demilew et al, 2018. Treatment 1 (R2 and R3) showed the starting

point was on day 9, followed by Treatment 1 (RS) and Treatment 2 (R1, R4, and RS) on day 10. Then, Treatment 3 (R1, R2, and RS) on day 13. It was followed by Treatment 1 (R1 and R4), Treatment 2 (R3) and Treatment 4 (R1 and R3) on day 14. In addition, Treatment 2 (R2) and Treatment 3 (R3) on day 15. Moreover, Treatment 3 (R4) on day 16. Lastly, black bars represent Treatment 4 (R2, R4, and RS), which did not undergo epithelization during the experimental phase.




Figure 3. Data on Epithelization Period.

Normal epithelization occurs within 14 days as seen in studies conducted by Ozeion LLC, Wilmington, DE - (2013) and the results of the study conform to it. Early epithelization of some replicates in Paragis extract and Paragis ointment may be due to the presence of secondary metabolites such as flavonoids, alkaloids, and tannins as cited by Iberahim ct al. (2015).




Figure 4. Data on Wound Healing.

Based on the gathered results, the graph shows that Treatment 1 (R2, R3, and RS) and Treatment 2 (R1, R4, and

RS) were fully healed on day 13. Treatment 1 (R1, and R4), Treatment 2 (R3), Treatment 3 (R1, R2 and RS) and Treatment 4 (R3) were fully healed on day 16. Moreover, black 'bars represent Treatment 2 (R2), Treatment 3 (R3 and R4) and Treatment 4 (R1, R2, R4 and RS) which did not heal during the experimental phase.

The normal fully healed wound occurs within 7-21 days as seen in studies conducted by - K. Kowalski et al. (2017) and the study results conform to it. The rapid healing of wounds seen in Paragis extract and Paragis ointment may be due to the presence of secondary metabolites like alkaloids, flavonoids, and tannins, as cited by Tberahim et al. (2015).

CONCLUSION

Based on the results, the researchers conclude that the Paragis leaf ethanolic extract and ointment have a burn wound healing property, with Treatment 1 being more effective than Treatment 2 due to the presence of an emulsifying agent that increases retention onto the skin of test animals. Test substance such as the ointment and ethanolic leaf extract was confirmed not toxic, for the test animals did not show severe and enduring signs of distress and pain.

RECOMMENDATION

The study on the burn wound healing potential of the ethanolic extract of Paragis (*E. indica*) leaves has yielded promising results, indicating the need for further exploration. To build upon these findings, several recommendations have been put forth. Firstly, it is advisable to extend the investigation to encompass other types of wounds, such as excision and incision wounds, to comprehensively assess the healing capabilities of Paragis Extract and its potential application in various clinical scenarios. Secondly, alternative methods of extraction should be explored to determine if different processes yield extracts with enhanced wound healing properties.


Moreover, additional physicochemical evaluations, including assessments of viscosity, spreadability, and solubility of the ointment containing Paragis Extract, can provide valuable insights into its formulation and practical application. Furthermore, to gain a more comprehensive understanding of its therapeutic effects, it is recommended to examine various wound healing parameters, including wound coagulation, inflammation, granulation, and maturation.

APPENDICES


Appendix A. Percent Wound Contraction


TREATMENTS/DAYS	1	4	7	10	13	16
OINTMENT	0	10.82	26.66	45.79	66.60	100
PARAGIS EXTRACT	0	11.66	24.99	40.81	77.48	93.32
NITROFURA ZONE	0	6.64	13.34	27.5	54.96	84.98
UNTREATED	0	2.49	9.99	16.68	26.66	66.62

Acute Dermal



Burn Wound Healing





REFERENCES

- Abdullahi, A., Amini-Nik, S., & Jeschke, M. G. (2014).** Animal models in burn research. *Cellular and Molecular Life Sciences*, 71(17):3241-3255. doi: 10.1007/500018-014-1612-5
- Ahmed, S., Yousaf, M., Mothana, R. A., & Al-Rehaily, A. (2016).** Studies on wound healing activity of some Euphorbia species on experimental rats. *African Journal of Traditional, Complementary, and Alternative Medicines*, 13(5), 145-152. doi: 10.21010/ajtcam.v13i5.19
- Alaekwe, I. O., Ajiwe, V. I. E., Ajiwe, A. C., & Aningo, G. N. (2015).** Phytochemical and antimicrobial screening of the aerial parts of *Eleusine indica*. *International Journal of Pure & Applied Bioscience*, 31), 257-264. Retrieved from ijfab website, <http://www.ijfab.com/form/2015%20Volume%203.%20issue%201/1JPAB-2015-3-1-257-264.pdf>
- Al-Zubairi, A. S. Abdul, A. B., Abdelwahab, S. I., Peng, C. Y., Mohan, S., & Elhassan, M. M. (2011).** *Eleusine indica* possesses antioxidant, antibacterial and cytotoxic properties. *Evidence-Based Complementary and Alternative Medicine*, 2011, 1-6. doi.org/10.1093/ecam/nep091
- Amoah, D. O., Joson, M. B., & Pareja, M. C. (2017).** Antiuro lithiatic potential of *Eleusine indica* Linn. (goose grass) root extract on ethylene glycol induced nephrolithiasis in *Rattus norvegicus* (albino rats). *Biomedical Sciences*, 3(5), 99-108. doi: 10.11648/j.bs.20170305.13
- Babu, R. H. & Savithamma, N. (2014).** Studies on mineral analysis of grasses of Poaceae. Retrieved from Semantic Scholar website, [aper/Studies-on-Mineral-Analysis-of-Grasses-of-Poaceae-Babu Savithamma/2cc57b272ec97ce](aper/Studies-on-Mineral-Analysis-of-Grasses-of-Poaceae-Babu-Savithamma/2cc57b272ec97ce) 3c7 d0671d20096f3c18fe6fbe
- Baravkar, A. A., Kale, R. N., Patil, R., & Sawant, S. D. (2008).** Pharmaceutical and biological evaluation of formulated cream of methanolic extract of *Acacia nilotica* leaves. *Research Journal of Pharmacy and Technology*, 1(4), 481-483. Retrieved from Research Gate website, <https://www.researchgate.net/publication/285320500-Pharmaceutical-and-biological-evaluation-of-formulated-cream-of-methanolic-extract-of-Acacia-nilotica-leaves>.
- Beecham, G. B., Bansal, P., Nessel, T. A., & Goyal, A. (2021, January 26).** Lidocaine. StatPearls Publishing. www.ncbi.nlm.nih.gov/books/NBK539881/

- Bhatia, N., Kaur, G., Soni, V., Kataria, J., & Dhawan, R. K. (2016).** Evaluation of the wound healing potential of isoquercetin-based cream on scald burn injury in rats. *Burns & Trauma*, 4(7), 1-8. doi:10.1186/s41038-016-0032-1
- Bhattacharya, S. (2012).** Wound healing through the ages. *Indian Journal of Plastic Surgery*, 45(2), 177. doi:10.4103/0970-0358.101255
- Cai, E. J., Ang, C. H., Raju, A., Tan, K. B., Hing, E. C. H., Loo, Y., & Lim, T. C. (2014).** Creation of consistent burn wounds: A rat model. *Archives of Plastic Surgery*, 41(4), 317-324. https://doi.org/10.5999/aps.2014.41.4.317
- De Oliveira, A. A., & Romo, N. F. (2015).** Growth inhibition and pro-apoptotic action of *Eleusine indica* (L.) Gaertn extracts in allium test. *European Journal of Medicinal Plants*, 8(3), 121-127. https://doi.org/10.9734/EJMP/2015/18099
- Demilew, W., Adinew, G. M., & Asrade, S. (2018).** Evaluation of the wound healing activity of the crude extract of leaves of *Acanthus polystachyus* Delile (Acanthaceae). *Evidence-Based Complementary and Alternative Medicine*, 2018, 1-9. https://doi.org/10.1155/2018/2047896
- Ettebong, E. O., Nwafor, P. A., Ubulom, P. E., & Udobang, J. A. (2017).** The toxicological assessment of ethanolic whole-plant extract of *Eleusine indica* in Wistar albino rats. *Journal of Herbmed Pharmacology*, 6(2), 49-54. http://herbmedpharmacol.com
- Ettebong, E. O., Etukakpan, E., & Bassey, A. (2016).** Anticonvulsant potentials of ethanolic extract of *Eleusine indica*. *Journal of Coastal Life Medicine*, 4(12), 965-968. <https://doi.org/10.12980/jclm.4.2016j6-196>
- Ettebong, E. O., & Bassey, A. I. (2015).** In vitro antimicrobial evaluation of whole-plant extracts of *Eleusine indica*. *Journal of Medicinal Plants Studies*, 5(4), 99-102. https://www.plantsjournal.com
- Gan, D., Su, Q., Su, H., Wu, L., Chen, J., Han, B., & Xiang, M. (2021).** Burn ointment promotes cutaneous wound healing by modulating the PI3K/AKT/mTOR signaling pathway. *Frontiers in Pharmacology*, 12, 1-13. https://doi.org/10.3389/fphar.2021.631102
- Garba, S., Osemeahon, S., Maina, H. M., & Barminas, E. (2012).** Ethylenediaminetetraacetate (EDTA)-assisted phytoremediation of heavy metal contaminated soil by *Eleusine indica* L. Gaertn. *Journal of Environmental Chemistry and Ecotoxicology*, 4, 103-109.
- Guevara, B. Q. (2005).** A guidebook to plant screening: Phytochemical and biological. University of Santo Tomas Publishing House.
- Guo, H.-F., Ali, R. M., Hamid, R. A., Zaini, A. A., & Khaza'ai, H. (2017).** A new model for studying deep partial-thickness burns in rats. *International Journal of Burns and Trauma*, 7(6), 107-114. https://pubmed.ncbi.nlm.nih.gov/29119063
- Heuzé, V., Tran, G., Hassoun, P., & Lebas, F. (2019, July 30).** Goosegrass (*Eleusine indica*). *Feedipedia*. https://www.feedipedia.org/node/446
- Hopf, H. W., Kelly, M., & Shapshak, D. (2008).** Oxygen and the basic mechanisms of wound healing. In *Physiology and medicine of hyperbaric oxygen therapy* (pp. 203-228). https://doi.org/10.1016/B978-1-4160-3406-3.50016-0
- Iberahim, R., Bahtiar, A., & Ibrahim, N. (2016).** Anti-herpes simplex virus type-1 activity of *Eleusine indica* methanol extract. *Malaysian Journal of Microbiology*, 12, 71-77.

- Iberahim, R., Yaacob, A., & Ibrahim, N. (2015).** Phytochemistry, cytotoxicity and antiviral activity of *Eleusine indica* (sambau). In *The 2015 UKM FST Postgraduate Colloquium: Proceedings of the Universiti Kebangsaan Malaysia*.
<https://doi.org/10.1063/1.4931234>
- Iqbal, M., & Gnanaraj, C. (2012).** *Eleusine indica* L. possesses antioxidant activity and precludes carbon tetrachloride (CCl₄)-mediated oxidative hepatic damage in rats. *Environmental Health and Preventive Medicine*, 17(4), 307-315.
<https://doi.org/10.1007/s12199-011-0255-5>
- Khoo, Y. T., Halim, A. S., Singh, K. K., & Mohamad, N. A. (2010).** Wound contraction effects and antibacterial properties of tualang honey on full-thickness burn wounds in rats in comparison to hydrofibre. *BMC Complementary and Alternative Medicine*, 10, 48.
<https://pubmed.ncbi.nlm.nih.gov/20815896>
- Masson-Meyers, D. S., Andrade, T. A. M., Caetano, G. F., Guimarães, F. R., Leite, M. N., Leite, S. N., & Frade, M. A. C. (2020).** Experimental models and methods for cutaneous wound healing assessment. *International Journal of Experimental Pathology*, 101, 21-37.
<https://doi.org/10.1111/iep.12346>
- Mayo Clinic. (n.d.).** Nitrofurazone (topical route). Retrieved May 27, 2021, from <https://www.mayoclinic.org/drugs-supplements/nitrofurazone-topical-route/side-effects/drg-20065093>
- Morah, F., & Otuk, M. E. (2015).** Antimicrobial and antihelmintic activity of *Eleusine indica*. *Acta Scientiae et Intellectus*, 1(4), 28-32.
- Nagar, H. K., Srivastava, A. K., Srivastava, R., Kurmi, M. L., Chandel, H. S., & Ranawat, M. S. (2016).** Pharmacological investigation of the wound healing activity of *Cestrum nocturnum* (L.) ointment in Wistar albino rats. *Journal of Pharmaceutics*, 2016, 1-8.
<https://doi.org/10.1155/2016/9249040>
- Nas, J., Dangeros, S., Chen, P., Dimapilis, R., Gonzales, D., & Hamja, F. (2020).** Evaluation of anticancer potential of *Eleusine indica* methanolic leaf extract through Ras- and Wnt-related pathways using transgenic *Caenorhabditis elegans* strains. *Journal of Pharmaceutical Negative Results*, 11(1), 42-46.
https://doi.org/10.4103/jpnr.JPNR_7_20
- National Center for Biotechnology Information. (2021a).** PubChem compound summary for CID 3676, lidocaine. Retrieved May 27, 2021, from <https://pubchem.ncbi.nlm.nih.gov/compound/Lidocaine>
- National Center for Biotechnology Information. (2021b).** PubChem compound summary for CID 5447130, nitrofurazone. Retrieved May 27, 2021, from <https://pubchem.ncbi.nlm.nih.gov/compound/Nitrofurazone>
- Okokon, J. E., Odomena, C. S., Effiong, I., & Obot, J. (2010).** Antiplasmodial and antidiabetic activities of *Eleusine indica*. *International Journal of Drug Development & Research*, 2(3), 493-500.
- Okokon, J. E., Udoh, A. E., Etebong, E., & Adamu, A. B. (2014).** Antioxidative burst, cytotoxic and antileishmanial activities of *Eleusine indica*. *International Journal of Clinical Pharmacology Research*, 4(3), 165-168.

- Ong, S. L., Mah, S. H., & Lai, H. Y. (2016). Porcine pancreatic lipase inhibitory agent isolated from medicinal herb and inhibition kinetics of extracts from *Eleusine indica* (L.) Gaertner. *Journal of Pharmaceutics*, 2016. <https://doi.org/10.1155/2016/8764274>
- Ong, S. L., Nalamolu, K. R., & Lai, H. Y. (2017). Potential lipid-lowering effects of *Eleusine indica* (L.) Gaertn. extract on high-fat-diet-induced hyperlipidemic rats. *Pharmacognosy Magazine*, 13(Suppl. 1), S81-S89. <https://doi.org/10.4103/0973-1296.203986>
- Organization for Economic Cooperation and Development. (2017). Test No. 402: Acute dermal toxicity. OECD Publishing. <https://doi.org/10.1787/9789264070585-en>
- Oyedeeji, S., Raimi, I. O., & Odiwe, A. L. (2013). A comparative assessment of the crude oil-remediating potential of *Cynodon dactylon* and *Eleusine indica*. *Environment Pollution Ecology*, 11, 145-150.
- Pastar, I., Stojadinovic, O., Yin, N. C., Ramirez, H., Nusbaum, A. G., Sawaya, A., & Tomic-Canic, M. (2014). Epithelialization in wound healing: A comprehensive review. *Advances in Wound Care*, 3(7), 445-464. <https://doi.org/10.1089/wound.2013.0473>
- Pattanayak, S., Nayak, S. S., Dinda, S. C., Panda, D., & Navale, K. P. (2011). Evaluation of herbal ointments formulated with methanolic extract of *Cajanus scarabaeoides*. *Journal of Pharmacy and Allied Health Sciences*, 1, 49-57. <https://doi.org/10.3923/jpahs.2011.49.57>
- Pereira, D. S. T., Lima-Ribeiro, M. H. M., de Pontes-Filho, N. T., Carneiro-Leão, A. M. A., & Correia, M. T. S. (2012). Development of animal model for studying deep second-degree thermal burns. *Journal of Biomedicine and Biotechnology*, 2012, 1-7. <https://doi.org/10.1155/2012/460841>
- Porumb, V., Trandabat, A. F., Terinte, C., Caruntu, I. D., Porumb-Andrese, E., Dimofte, M. G., & Pieptu, D. (2017). Design and testing of an experimental steam-induced burn model in rats. *BioMed Research International*, 2017, 1-10. <https://doi.org/10.1155/2017/9878109>
- Sagnia, B., Fedeli, D., Casetti, R., Montesano, C., Falcioni, G., & Colizzi, V. (2014). Antioxidant and anti-inflammatory activities of extracts from *Cassia alata*, *Eleusine indica*, *Eremomastax speciosa*, *Carica papaya*, and *Polyscias fulva* medicinal plants collected in Cameroon. *PLoS ONE*, 9(8), 1-10. <https://doi.org/10.1371/journal.pone.0103999>
- Shedoeva, A., Leavesley, D., Upton, Z., & Fan, C. (2019). Wound healing and the use of medicinal plants. *Evidence-Based Complementary and Alternative Medicine*, 2019, 1-25. <https://doi.org/10.1155/2019/2684108>
- Stuart, G. U. (2013). *Paragis: Eleusine indica* (Linn.) Gaertn. Retrieved May 27, 2021, from <http://stuartxchange.org/Paragis>
- Subalakshmi, M., Saranya, A., Maheswari, M. U., Jarina, A., & Kavimani, S. (2014). An overview of the current methodologies used for the evaluation of drugs having wound healing activity. *International Journal of Experimental Pharmacology*, 4, 127-131.

Thakur, R., Jain, N., Pathak, R., & Sandhu, S. S. (2011).

Review article: Practices in wound healing studies of plants. Evidence-Based Complementary and Alternative Medicine, 2011, 1–17.
<https://doi.org/10.1155/2011/438056>

Tutor, J. T., & Chichioco-Hernandez, C. L. (2018).

Angiotensin-converting enzyme inhibition of fractions from *Eleusine indica* leaf extracts. Pharmacognosy Journal, 10(1), 25–28.
<https://www.phcogj.com/sites/default/files/PharmacognJ-10-25.pdf>

Wallace, H. A., Basehore, B. M., & Zito, P. M. (2020, September 29). Wound healing phases. StatPearls Publishing.

<https://www.ncbi.nlm.nih.gov/books/NBK470443/>

Calentig, P. G. T. (2019). Phytochemical screening and antibacterial activity of *Eleusine indica* (Paragis) flower and leaf extract.
<https://ojs.aaresearchindex.com>

ACKNOWLEDGEMENT

We would like to extend our sincere acknowledgments to all those who played a pivotal role in completing this study on the burn wound healing potential of Paragis (*Eleusine indica*) leaf ethanolic extract. Firstly, we express our heartfelt gratitude to our dedicated research team, whose unwavering commitment and collective efforts were instrumental in carrying out this research.

We also thank Cagayan State University, Carig Campus, for its support in providing laboratory facilities and equipment that enabled us to conduct our research with precision and dedication.

Note from the author: The accuracy and integrity of the content in this article are the sole responsibility of the author(s).