




Water Buffalo as a Reservoir for *Schistosoma Japonicum*: Policy Implications for Animal and Human Health



Public Health and Medicine

Vincent Jim P. Palor, DVM*

Cagayan State University, College of Agriculture, Gonzaga Campus, Gonzaga, Philippines

Corresponding Author: vincentjimpalor@csu.edu.ph

Key Findings and Policy Implications

- A recent study in Gonzaga, Cagayan, Philippines found that water buffaloes in 14 barangays had high rates of *Schistosoma japonicum* (22.60%) and *Fasciola* (83.68%) infections. Only eleven barangays reported *S.*, although all had *Fasciola* cases. *japonicum*, with the largest risk of infection and pollution found in Tapel and Magrafil. Severe *S.* was substantially more common in older animals. *japonicum* infections; however, there was no discernible sex-based difference.
- Crucially, co-infection was widespread, and there was a strong association between *S. japonicum* and *Fasciola*.

presence of, indicating that animals with one parasite infection are more vulnerable to the other. To safeguard animal welfare and reduce the danger of zoonotic transmission, these findings emphasize the necessity of focused veterinary interventions, farmer education, and enhanced parasite control programs.

- Stronger animal health surveillance and parasite management initiatives are urgently needed in endemic locations, as evidenced by the high frequency of *Schistosoma japonicum* and *Fasciola* infections among water buffaloes in Gonzaga, Cagayan. The necessity of

regular veterinary deworming and focused treatment approaches is highlighted by the strong correlation between co-infection and increased illness severity, especially in older animals.

- Integrating livestock management within the Department of Health's current *schistosomiasis* control initiatives should be a top priority for policymakers. This involves community education on parasite transmission and safe livestock handling, frequent fecal surveillance, and veterinarian services that are subsidized. Putting money into these preventative measures can boost farm output, lessen the burden of disease in both humans and animals, and aid in the larger objective of zoonotic disease control in rural agricultural communities.

Background

In the Philippines, *schistosomiasis* is still a major agricultural and public health issue, especially in rural areas where livestock are essential to farming. Humans and animals can contract the parasite trematode *Schistosoma japonicum* by coming into touch with water sources that are home to infected snails. Carabaos, also known as water buffaloes, are frequently employed in rice cultivation and are important reservoir hosts that support the parasite's ongoing spread.

Because of the possible effects on both animal productivity and public health, local stakeholders in Gonzaga, Cagayan, are concerned about reports of probable illnesses in carabaos. Water buffaloes are especially susceptible to parasite illnesses, such as *S*, because to their wallowing nature and ongoing contact with rice paddies and irrigation canals. *Fasciola* and *japonicum*.

Designing efficient control strategies for water buffaloes requires an understanding of the prevalence and risk factors of these illnesses. The purpose of this study was to determine the degree of *Schistosoma japonicum* infection in water buffaloes in a few Gonzaga barangays and to look for potential correlations with age, sex, and co-infection with *Fasciola*. In order to lower infection risks, protect animal welfare, and slow the spread of zoonotic illnesses in the area, the findings are meant to serve as a guide for local farmers, veterinarians, and legislators as they adopt focused interventions.

Research Objectives

This study aimed finding the prevalence of *Schistosoma japonicum* infection in water buffaloes in a few Gonzaga, Cagayan barangays and identifying risk variables including age, sex, and co-infection with *Fasciola* were the objectives of this study. In order to lower the danger of zoonotic transmission and decrease parasite illnesses in livestock, the results aim to support improved control measures.

Methodology

Experimental design

The experiment employed a descriptive design with correlation analysis. The study's significance was also evaluated using the T-test. The frequency of *S. Schistosome* eggs found in fecal samples were used to diagnose *japonicum* infection in water buffaloes. Five grams of excrement were counted for eggs. To determine its relevance in relation to the occurrence of *S*, the animals' ages were also taken into account. infection with *japonicum*. Pearson Product-Moment Correlation analysis (Pearson rho) and the associated t-test of the correlation's significance were used to ascertain this. Using a t-test for two independent samples, the association between sex and prevalence was ascertained by comparing the prevalence of male and female animals. The statistical significance was determined at a 0.05 level.

In particular, the total number of eggs in 10 g of fecal samples was used to calculate the degree of infection among water buffaloes in each endemic area. Additionally, according to Ardina (2014), the number of eggs in 10 g of feces was used to evaluate the strength of infection, which was then classified as light (1-400 eggs), moderate (101-400 eggs), or heavy (>400 eggs). Each animal's bovine contamination index (BCI) was calculated.

Key Findings




Figure 1. Ranking of affected Barangay with *Schistosoma japonicum*.




Figure 2. Ranking of affected Barangay with *Fasciola*.

Table 1. Number and percentage of eggs per sample.

Variables	Total	Mean
Total Number of Treatments	478	
Total Eggs pre gram (<i>Fasciola</i>)	16,920	35.4
Total Eggs pre gram (<i>S. japonicum</i>)	2,265	4.74

Table 2. Number of animals affected by the parasite.

Parasite	Number of Animals Affected	Percentage
Number if animals with <i>Fasciola</i>	400	83.68%
Number if animals with <i>S. japonicum</i>	109	22.80%
Total	509	106.49%

Table 1. shows that out of 478 water buffalo tested *Fasciola* had a total of 16,920 eggs with a mean of 35.4, while the *S. japonicum* had a total of 2,265 eggs with a mean of 4.74.

Table 2. shows that out of the 478 samples, 400 were infected with *Fasciola*, which constitutes 83.68%. There were 109 heads infected with *S. japonicum*, which is 22.60% of the animals tested. Of the total animals infected, there were those who were infected with *Fasciola* alone, others were infected with *S. japonicum* alone, and still, others were infected by both parasites.

Conclusions

According to the study, *Schistosoma japonicum* is common in water buffalo, indicating that these animals have a role in the parasite's transmission in human-infecting water bodies. There will be a significant decrease in the quantity of parasites in the Gonzaga area that is afflicted if the proper steps are taken to limit the animals' accessibility to water bodies and to conduct widespread deworming.

Recommendations

- To prevent *Schistosoma japonicum* and *Fasciola* infections, it is advised that local government entities, in collaboration with the Department of Agriculture and the Department of Health, conduct routine, community-wide water buffalo deworming programs. Treatment plans should give priority to older animals because they display higher infection levels.
- Additionally, barangays like Magrafil, Tapel, and Sta. Cruz of Gonzaga, Cagayan, Philippines have the highest rates of pollution and infection. should be prioritized for focused treatments, such as fecal screening, snail management, and restricting buffalo access to water

bodies and irrigation canals, which are known to be transmission sources.

- Support should also be given to initiatives aimed at increasing capacity, such as educating farmers on environmental sanitation, animal husbandry, and parasite prevention. Including livestock parasite control in schistosomiasis prevention programs will lower the danger of zoonotic transmission to humans while simultaneously safeguarding farm productivity and animal health.

Target Policy Actors and Beneficiaries



References

- Navarro, J. R. P., Sunico, L. S., Martinez, R. M., Sison, S. M., Esparar, D. G., Villar, L. C., Belizario, V. Y., Jr., Velasco, L. R., & de Leon, W. U. (2005). Cagayan Valley: A newly described endemic focus for Schistosomiasis japonicum in the Philippines. *Philippine Journal of Internal Medicine*, 43(3), 117-122.
- Dumag, P. U., Gajudo, C. E., Sena, C. Y., Cardenas, E. C., & Fementira, E. B. (1981). Epidemiology of animal schistosomiasis in the Philippines. *The Philippine Journal of Animal Industry*, 36, 1-23.
- Gordon, C. A., Acosta, L. P., Gray, D. J., Olveda, R. M., Jarilla, B., McManus, D. P., Williams, G. M., Rivera, P. T., & Ross, A. G. (2012). High prevalence of *Schistosoma japonicum* infection in carabao from Samar Province, the Philippines: Implications for transmission and control. *PLOS Neglected Tropical Diseases*, 6(9), Article e1778. <https://doi.org/10.1371/journal.pntd.0001778>
- Guo, J., Li, Y., Gray, D., Ning, A., Hu, G., Chen, H., Davis, G. M., Sleight, A. C., Feng, Z., & McManus, D. P. (2006). A drug-based intervention study on the importance of buffaloes for human *Schistosoma japonicum* infection around Poyang Lake, People's Republic of China. *American Journal of Tropical Medicine and Hygiene*, 74(2), 335-341.
- Editor's Note**
- This policy brief is based on results of the CSU-funded project "Water Buffalo as a Reservoir for *Schistosoma japonicum*: Policy Implications for Animal and Human Health" This project was implemented by the Cagayan State University - Gonzaga Campus. The CSU - Gonzaga Campus project is composed of Dr. Vincent Jim P. Palor.

About CSU Policy Brief

Our policy brief offers an in-depth analysis of contemporary challenges and opportunities within higher education, with a special focus on the unique context of Cagayan State University. Through rigorous research and stakeholder engagement, we examine key areas such as academic innovation, research advancement, community engagement, and administrative excellence. Each recommendation is carefully developed to foster an environment of growth, equity, and sustainability. **This policy brief is published bi-annually.**

Our approach is built on transparency and inclusiveness. By integrating comprehensive data analysis with practical insights from experts and the broader university community, we ensure that our policy recommendations are not only academically robust but also practically implementable. This process underscores our commitment to creating a positive and lasting impact on the university's future.

The policy brief also serves as a platform for translating research outputs into practical policy insights across six strategic thematic areas including Science and Technology in Agriculture, Natural Resources, Education, Food Science and Chemistry, Public Health and Medicine, and Behavioral and Social Sciences. These areas reflect the university's research priorities and address emerging concerns related to agriculture and livestock, environmental management, teaching and learning, food innovation, clinical diagnostics, and community resilience.

While these publications provide science-based interventions for regional and national development, the analyses and recommendations are those of the authors and do not necessarily represent the official positions of Cagayan State University. Any reproduction, distribution, or citation of this work must acknowledge the author and the CSU Policy Brief.

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means without prior permission from Cagayan State University. This Policy Brief is part of the CSU Policy Brief.



Get in touch with us via:

-  facebook.com/csuktmpo
-  csu_ktmooffice@csu.edu.ph
-  csu.org.ph/policybrief



Cagayan State University

Knowledge & Technology Management and Publication Office
RDE Bldg., CSU- Andrews Campus Caritan Sur, Tuguegarao City, Cagayan

Editorial Board

Gilbert C. Magulod Jr., PhD
Editor-in-Chief

Patricianne M. Padua, PhD
Associate Editor-in-Chief

Peer Reviewers

Junel B. Guzman, PhD
Josie Y. Bas-Ong, PhD
Danica B. Ibañez
Arianne Faye D. Perucho, MD

Editorial Staff

Jamaica C. Pedro
Managing Editor

Marcel W. Visitacion
Layout Artist

Technical Staff

Christian Dave E. Columna
IT Technician

Interagency Local Advisory Board

Florentina S. Dumlao, PhD
University Consultant, Cagayan State University

Maria Mercedes Arzadon, PhD
University of the Philippines

Atty. Norman Clarence T. Lasam
Cagayan State University

Engr. Gil Mark S. Hizon
Department of Science and Technology Region II

Emma L. Ballad, PhD
Bureau of Fisheries and Aquatic Resources Region II

Arthur G. Ibañez, PhD, ASEAN Engr.
OIC, Office of the University President



POLICY BRIEF

Cagayan State University
Knowledge & Technology Management and Publication Office



csu.org.ph/policybrief